

In this Guideline:

- [Referral to Rule Out Cataract](#)
- [Protocol for Diagnosis of Congenital Cataracts in Children](#)
- [References](#)


Referral to Rule Out Cataract

- Children **under 8 weeks old** with suspected cataract need to be seen in the next 48 hours
- Children **from 8 weeks old to 8 years old** should be seen within 2 weeks
- Children **over 8 years old** should be seen within 4 weeks

Proceed to [protocol for diagnosis of congenital cataracts](#).

Protocol for Diagnosis of Congenital Cataracts in Children

Return to Page 1



References



1. BCBS 2011-2012
2. Taylor, D. Congenital Cataract: The History and The Practice. *Journal of Eye*. 1998, 12 9-36
3. *Ophthalmology Clinical Signs and Differential Diagnosis*. Jack J Kanski, Ken K Nischal.
4. *The Wills Eye Manual. Office and Emergency Room. Diagnosis and Treatment of Eye Disease*. Fifth edition.
5. Birch E et al. *Early Treatment of Congenital Unilateral Cataract Minimizes Unequal Competition*. IOVS
6. Graw J. Congenital Hereditary Cataract. *Int J Dev Biol*. 48: 1031-1044.
7. UpToDate
8. Fledelius, Christensen. Reappraisal of human ocular growth curve. *BJO* 96;80:918
9. Mutti et al. Axial growth and changes in lenticular and corneal power during emmetropization. *IOVS* 05; 46: 3074
10. Loyd IC, Ashworth J, Biswas S, Abadi RV. Advances in the management of congenital and infantile cataract. *Eye (London)* 2007. Oct; 21(10):1301-9
11. Hosal & Biglan. Risk factors for secondary membrane formation after removal of paediatric cataract. *JCRS* 02; 28:302
12. Basti et al. Results of a prospective evaluation of three methods of management of paediatric cataract. *Ophthalmology* 96;103:713
13. Vasavada et al. Primary posterior capsulorhexis with and without anterior vitrectomy in congenital cataracts. *JCRS* 97;23;645
14. Koch & Kohen. Retrospective comparison of techniques to prevent secondary cataract formation after posterior chamber IOL in infants and children. *JCRS* 97;23:657
15. Ben Ezra et al. Posterior capsulectomy in paediatric cataract surgery; the necessity of a choice. *Ophthalmology* 97; 104:2168.
16. Lloyd, Kriss, Speedwell, Thompson, Russell-Eggitt et al. Modulation of Amblyopia therapy following early surgery for unilateral congenital cataracts. *Br. J. Ophthalmol*. 1995;79;802-806.
17. The Infant Aphakia Treatment Study Group. The infant Aphakia Treatment Study: Design and Clinical Measures at Enrollment. *Arch Ophthalmol*. 128(1):N21-27.2010.

Guideline Developed by: Ken Nischal, MD; Matthew Pihlblad, MD; Preeti Patil, MD; Amgad Eldib, MD; Jamila Hiasat, MD; Marie-Helene Errera, MD; Erin Keim, OD; Craig Luchansky, OD; Ellen Mitchell, MD; Elizabeth Conner, MD; Meghal Gragani, MD; Alki Liasis, BSc, PhD, Hannah Scanga, MS, CGC; Kelly Schooping Tripi, MS, CGC; Kaajal Nanda, BMedSci

#: 807.00

Origination: 05/2016

Revised: 08/2021

*A Clinical Effectiveness Guideline (CEG) is an evidence-based, **suggested** collaborative care plan that is intended to promote coordination and communication with respect to patient care and may be modified to meet individual care needs.*